


Stabilis



Nystatin



Stabilité des préparations

		2.29 MUI Mycostatine®	NaHCO3 1.4% >> 500 ml	19-23°C		4		2820
		2.29 MUI Mycostatine®	NaHCO3 1.4% >> 500 ml	19-23°C		4		2820
		2.29 MUI Mycostatine®	NaHCO3 1.4% >> 500 ml	4°C		15		2820
		2.4 MUI = 24 ml Mycostatine®	Hexetidine 0.1% 75 ml NaHCO3 1.4% >>400 ml	20-25°C		3		2729
























Bibliographie

	Type	Source
2729	Revue	Vincent I, Schmit B, Houze S, benhaïem N, Jacques L, Taburet AM. Bains de bouche antifongiques: études de stabilité. J Pharm Clin 1995 ; 14, 2: 106-111.
2820	Revue	Groeschke J, Solassol I, Bressolle F, Pinguet F. Stability of amphotericin B and nystatin in antifungal mouthrinses containing sodium hydrogen carbonate J Pharm Biomed Anal 2006 ; 42: 362-366.



Dictionnaire

 Antifongique	 Gargarisme
 Stabilité des préparations	 Contenant
 Origine	 Excipient
 Température	 Conservation
 Durée de stabilité	 Biosimilaire
 Données conflictuelles	 Bibliographie
 Verre	 Poudre
 A l'abri de la lumière	 Jour
 Polypropylène	 Non précisé
 Liquide	 Bibliographie
 Dictionnaire	