


Stabilis



Rifabutin



Stabilité des préparations

		200 mg Mycobutin®	Sirop de cerise >>10ml	25°C	?	84		2419
		200 mg Mycobutin®	OraPlus® / OraSweet® (1:1) >>10ml	25°C	?	84		2419
		200 mg Mycobutin®	Sirop de cerise >>10ml	30-40°C	?	56		2419
		200 mg Mycobutin®	OraPlus® / OraSweet® (1:1) >>10ml	30-40°C	?	84		2419
		200 mg Mycobutin®	Sirop de cerise >>10ml	4°C	?	84		2419
		200 mg Mycobutin®	OraPlus® / OraSweet® (1:1) >>10ml	4°C	?	84		2419




















Bibliographie

Type	Source
2419 Revue	Haslam JL, Egodage KL, Chen Y, Rajewski RA, Stella V. Stability of rifabutin in two extemporaneously compounded oral liquids Am J Health-Syst Pharm 1999 ; 56: 333-336.



Dictionnaire

 Antibiotique	 Solution buvable
 Stabilité des préparations	 Contenant
 Origine	 Excipient
 Température	 Conservation
 Durée de stabilité	 Biosimilaire
 Données conflictuelles	 Bibliographie
 Flacon plastique	 Gélules
 Non précisée	 Jour
 Bibliographie	 Dictionnaire